



**Super Cyclonic Storm “AMPHAN” over the southeast Bay of Bengal  
(16<sup>th</sup> – 21<sup>st</sup> May 2020): Summary**

**HIGHLIGHTS**

- The Super Cyclonic Storm (SuCS) “AMPHAN” (pronounced as UM-PUN) was the first SuCS over the BoB, after the Odisha SuCS of **1999**.
- It originated from the remnant of a Low Pressure Area which occurred in the near Equatorial Easterly wave over south Andaman Sea and adjoining southeast Bay of Bengal (BoB) on 13<sup>th</sup> May.
- It concentrated into a depression (D) over southeast BoB in the early morning of 16<sup>th</sup> May and further intensified into a deep depression (DD) in the same afternoon.
- It moved north- northwestwards and intensified into Cyclonic Storm over southeast BoB in the evening of 16<sup>th</sup> May, 2020. Moving nearly northwards, it further intensified into a Severe Cyclonic Storm (SCS) over southeast BoB in the morning of 17<sup>th</sup> May.
- It underwent rapid intensification during subsequent 24 hours and accordingly intensified into a Very Severe Cyclonic Storm (VSCS) by the afternoon of 17<sup>th</sup>, Extremely Severe Cyclonic Storm (ESCS) in the early hours of 18<sup>th</sup> and into a SuCS around noon of 18<sup>th</sup> May, 2020.
- It maintained the intensity of SuCS over westcentral BoB for nearly 24 hours, before weakening into an ESCS over westcentral BoB around noon of 19<sup>th</sup> May.
- Thereafter, it weakened slightly and crossed West Bengal – Bangladesh coasts as a VSCS, across Sundarbans, during 1530-1730 hrs IST of 20<sup>th</sup> May, with maximum sustained wind speed of 155 – 165 kmph gusting to 185 kmph. It lay over West Bengal as a VSCS, gradually moving north-northeastwards during late evening to night of 20<sup>th</sup> May. It moved very close to Kolkata during this period.
- Moving further north-northeastwards, it weakened into an SCS over Bangladesh & adjoining West Bengal around mid-night of 20<sup>th</sup> May, weakened further into a CS over Bangladesh in the early hours of 21<sup>st</sup> May, into DD over Bangladesh around noon of 21<sup>st</sup> May and into a D over north Bangladesh in the evening of the same day. It further weakened and lay as a well marked low pressure area over north Bangladesh and neighbourhood around mid-night of 21<sup>st</sup> May.
- **Monitoring of AMPHAN:**

India Meteorological Department (IMD) maintained round the clock watch over the north Indian Ocean and the system was monitored since 23<sup>rd</sup> April about three weeks prior to the formation of the Low Pressure Area on 13<sup>th</sup> May. In the extended range

outlook issued on 7<sup>th</sup> May, IMD indicated possible cyclogenesis during the second week over south Andaman Sea and adjoining southeast Bay of Bengal. On 9<sup>th</sup> May, it was indicated that a Low Pressure Area would form over the region on 13<sup>th</sup> May (96 hours prior to formation of the system) under the influence of the remnant cyclonic circulation persisting over the region during 6<sup>th</sup>-12<sup>th</sup>. On 11<sup>th</sup>, it was indicated that cyclogenesis (formation of depression) would occur around 16<sup>th</sup> May (48 hours prior to formation of the Low Pressure Area and 120 hours prior to formation of depression) over the BoB.

The cyclone was monitored with the help of available satellite observations from INSAT 3D and 3DR, polar orbiting satellites including SCATSAT, ASCAT etc. and available ships & buoy observations in the region. From 18<sup>th</sup> May midnight (1800 UTC) onwards till 20<sup>th</sup> May, the system was tracked gradually by IMD Doppler Weather Radars (DWRs) at Visakhapatnam, Gopalpur, Paradip, Kolkata and Agartala as it moved from south to north. IMD also utilised DWR products from 'DRDO Integrated Test Range', Chandipur, Balasore for tracking the system. Various numerical weather prediction models run by Ministry of Earth Sciences (MoES) institutions (viz.,IMD, IITM, NCMRWF & INCOIS), various global models and IMD's dynamical-statistical models developed in-house were utilized to predict the genesis, track, landfall and intensity of the cyclone. A digitized forecasting system of IMD was utilized for analysis and comparison of guidance from various models, decision making process and warning product generation.

➤ **Forecast Performance:**

**i) Genesis Forecast**

- The system was monitored since 23<sup>rd</sup> April about three weeks prior to the formation of Low Pressure Area over the southeast BoB on 13<sup>th</sup> May.
- In the extended range outlook issued on 7<sup>th</sup> May, cyclogenesis (formation of Depression) was predicted with low probability in the later part of week during 8<sup>th</sup>-14<sup>th</sup> May 2020. It was also predicted that the system would intensify further and move initially north-northwestwards and recurve north-northeastwards thereafter towards north BoB.
- In the Tropical Weather Outlook issued on 9<sup>th</sup> May, it was indicated that a Low Pressure Area would form over the region on 13<sup>th</sup> May (96 hours prior to formation of the system) under the influence of the remnant cyclonic circulation persisting over the region during 6<sup>th</sup>-12<sup>th</sup>.
- In the Tropical Weather Outlook issued on 11<sup>th</sup> May, it was indicated that cyclogenesis (formation of depression) would occur around 16<sup>th</sup> May (48 hours prior to formation of the Low Pressure Area and 120 hours prior to formation of depression) over the BoB. The Low pressure area formed on 13<sup>th</sup> May and concentrated into a Depression on 16<sup>th</sup> May morning.

**ii)Track, Intensity and Landfall Forecast**

- First information was provided in the **extended range outlook issued on 7th May (about 6 days prior to formation of LPA, 9 days prior to formation of**

**depression and 13 days prior to Landfall)** indicated that the system would intensify into a cyclonic storm and move initially northwestwards and recurve north-northeastwards towards north BoB

- In the **Tropical Weather Outlook, Press release and informatory message to the Government of India issued on 13th April (on the day of development of LPA, 3 days prior to formation of depression and 7 days prior to Landfall)**, it was indicated that the system would intensify into a cyclonic storm by 16th evening and would move initially northwestwards till 17th and then recurve north-northeastwards towards north BoB.
- Actually, the depression formed in the morning (0000 UTC) of 16th, cyclonic storm in the evening (1200 UTC) of 16th and the system moved north-northwestwards till 17th evening (1200 UTC) followed by north-northeastward recurvature thereafter and crossed West Bengal coast on 20th Afternoon.
- **In the first bulletin issued at 0845 IST of 16<sup>th</sup> May (104 hrs prior to landfall)** with the formation of Depression, it was indicated that the system would intensify into a cyclonic storm and will move north-northwestwards till 17<sup>th</sup> May followed by north-northeastward re-curved towards West Bengal coast during 18<sup>th</sup>-20<sup>th</sup> May and cross West Bengal coast with maximum sustained wind speed of 155-165 kmph gusting to 180 kmph.
- **In the bulletin issued at 1645 hrs IST of 16<sup>th</sup> May (24 hrs prior to rapid intensification)**, rapid intensification of the system was predicted and the system rapidly intensified from 17<sup>th</sup> afternoon onwards.
- **In the bulletin issued at 0845 hrs IST of 17<sup>th</sup> May (80 hrs prior to landfall)**, it was precisely mentioned that the system would cross West Bengal-Bangladesh coasts between Sagar Island (West Bengal) and Hatiya Islands (Bangladesh coast) during afternoon to evening of 20<sup>th</sup> May with maximum sustained wind speed of 155-165 kmph gusting to 185 kmph. The predicted track indicated Landfall across Sunderbans on 20<sup>th</sup> Afternoon.
- IMD continuously predicted since 16<sup>th</sup> May that Amphan will cross West Bengal coast as a very severe cyclonic storm (VSCS) with wind speed of 155-165 kmph gusting to 180 kmph on 20<sup>th</sup> May.

➤ **Cyclone warnings**

- In the first bulletin released at 0845 hrs IST of 16<sup>th</sup> May **(104 hrs prior to landfall), Pre-cyclone Watch** for West Bengal-north Odisha coasts was issued.
- The warnings were further upgraded and **Cyclone Watch** for West Bengal and north Odisha coasts was issued at 2030 hrs IST of 16<sup>th</sup> May **(92 hrs prior to landfall)**.
- **Cyclone Alert (Yellow Message)** for West Bengal and north Odisha coasts

was issued at 0840 hrs IST of 17<sup>th</sup> May **(80 hrs prior to landfall)**.

- **Cyclone Warning (Orange Message)** for West Bengal and north Odisha coasts was issued at 0845 hrs IST of 18<sup>th</sup> May **(56 hrs prior to landfall)**.
- **Post landfall outlook (Red Message)** for interior districts of Gangetic West Bengal, Assam and Meghalaya was issued at 2330 hrs IST of 19<sup>th</sup> May **(17 hrs prior to landfall)**.
- **Thus IMD predicted accurately the landfall point and time, track and intensity as well as associated adverse weather like wind, rainfall and storm surge due to cyclone, Amphan. We also predicted 100-110 gusting to 120 kmph wind speed and associated adverse weather for north Odisha coast, well in advance.**

### 1. Brief Life History:

- The Super Cyclonic Storm (SuCS) “AMPHAN” originated from the remnant of a Low Pressure Area which occurred in the near Equatorial Easterly wave over south Andaman Sea and adjoining southeast Bay of Bengal (BoB) during 1<sup>st</sup> – 5<sup>th</sup> May. Though the Low Pressure Area became less marked on 6<sup>th</sup> May, its remnant circulation meandered over south Andaman Sea and adjoining southeast BoB during 6<sup>th</sup> – 12<sup>th</sup> May. Under its influence, a fresh Low Pressure Area formed over southeast BoB and adjoining south Andaman Sea in the morning (0300 UTC) of 13<sup>th</sup> May.
- It lay as a well marked low pressure area (WML) over southeast BoB & neighbourhood in the morning (0300 UTC) of 14<sup>th</sup> May.
- Under favourable environmental conditions, it concentrated into a depression (D) over southeast BoB in the early morning (0000 UTC) of 16<sup>th</sup> May and further intensified into a deep depression (DD) in the same afternoon (0900 UTC).
- It moved north- northwestwards and intensified into Cyclonic Storm “AMPHAN” (pronounced as UM-PUN) over southeast BoB in the evening (1200 UTC) of 16<sup>th</sup> May, 2020. Moving nearly northwards, it further intensified into a Severe Cyclonic Storm (SCS) over southeast BoB in the morning (0300 UTC) of 17<sup>th</sup> May.
- It underwent rapid intensification during subsequent 24 hours and accordingly intensified into a Very Severe Cyclonic Storm (VSCS) by the afternoon (0900 UTC) of 17<sup>th</sup>, Extremely Severe Cyclonic Storm (ESCS) in the early hours of 18<sup>th</sup> (2100 UTC of 17<sup>th</sup> May) and into a Super Cyclonic Storm (SuCS) around noon (0600 UTC) of 18<sup>th</sup> May, 2020.
- It maintained the intensity of SuCS over westcentral BoB for nearly 24 hours, before weakening into an ESCS over westcentral BoB around noon (0600 UTC) of 19<sup>th</sup> May.

- Thereafter, it weakened slightly and crossed West Bengal – Bangladesh coasts as a VSCS, across Sundarbans, near latitude 21.65°N and longitude 88.3°E during 1530-1730 hrs IST (1000-1200 UTC) of 20<sup>th</sup> May, with maximum sustained wind speed of 155 – 165 kmph gusting to 185 kmph. It lay over West Bengal as a VSCS, gradually moving north-northeastwards during late evening to night (1200 – 1500 UTC) of 20<sup>th</sup> May. It moved very close to Kolkata during this period.
- Moving further north-northeastwards, it weakened into an SCS over Bangladesh & adjoining West Bengal around mid-night (1800 UTC) of 20<sup>th</sup> May, weakened further into a CS over Bangladesh in the early hours (2100 UTC of 20<sup>th</sup>) of 21<sup>st</sup> May, into DD over Bangladesh around noon of 21<sup>st</sup> May and into a D over north Bangladesh in the evening (1200 UTC) of the same day. It further weakened and lay as a well marked low pressure area over north Bangladesh and neighbourhood around mid night (1800 UTC) of 21<sup>st</sup> May.
- The observed track of the system during 16<sup>th</sup> – 21<sup>st</sup> May is presented in **Fig.1**. Best Track parameters associated with the system are presented in **Table1**.


## 2. Salient Features:

The salient features of the system were as follows:


- i. It was the first SuCS over the BoB, after the Odisha SuCS of 1999.
- ii. Climatologically, during satellite era a total of 7 severe cyclonic storms and above intensity storms developed within the grid  $\pm 2.5^\circ$  of the genesis point (10.4°N and 87.0°E). Out of these, 5 crossed Bangladesh coast and 2 crossed north Andhra Pradesh coast (Fig. 2a). Considering the total number of severe cyclonic storms and above intensity storms crossing West Bengal-Bangladesh coasts in the month of May during satellite era, a total of 2 (1989, 2009) severe cyclonic storms and above intensity storms crossed West Bengal coast and 7 crossed Bangladesh coast (Fig.2b).
- iii. It had a clockwise recurving track as it moved initially north-northwestwards till 1200 UTC of 17<sup>th</sup> and north-northeastwards thereafter. The total track length of the system was 1765 km. It was mainly steered by an anticyclonic circulation in middle & upper tropospheric levels to the northeast of the system centre.
- iv. It underwent intensification during 17<sup>th</sup> noon (0600 UTC) to 19<sup>th</sup> May early morning hours (2100 UTC of 18<sup>th</sup>) over westcentral BoB mainly due to low vertical wind shear (10-15 knots), very warm sea surface temperatures (SSTs – 30-31°C), high tropical cyclone heat potential (100 - 120 KJ /cm<sup>2</sup>) and increased cross equatorial wind surge. During this period, the system experienced an increase in maximum sustained wind speed (MSW) from 50 knots at 0600 UTC of 17<sup>th</sup> to 130 knots at 2100 UTC of 18<sup>th</sup> May.
- v. The peak MSW of the cyclone was 240-250 kmph (130 knots) gusting to 275 kmph (145 knots) during 1800 UTC of 18<sup>th</sup> to 0000 UTC of 19<sup>th</sup> May over the westcentral BoB. The lowest estimated central pressure was 920 hPa during

the same period (Fig.3a). Thereafter, the system started weakening over westcentral BoB under unfavourable environment (increase in vertical wind shear (20-25 knots)) and low Ocean thermal energy.


- vi. The system crossed West Bengal-Bangladesh coasts as a very severe cyclonic storm across Sundarbans, near lat.21.65°N/long. 88.30°E during 1000-1200 UTC, with maximum sustained wind speed of 85 knots gusting to 100 knots.
- vii. The system maintained the cyclonic storm intensity for almost 15 hours even after landfall from 1200 UTC of 20<sup>th</sup> May to 0300 UTC of 21<sup>st</sup> May.
- viii. The life period (D to D) of the system was 138 hours (5 days & 18 hours) against Long Period Average (LPA) of 134 hours (5 days & 14 hrs) for VSCS/ESCS categories over BoB during pre monsoon season based on the data of 1990-2013.
- ix. It moved with 12 hour average translational speed of 13.9 kmph against LPA (1990-2013) of 14.7 kmph for VSCS category over the north Indian Ocean (Fig.3b). During initial stages of it's development (0000 UTC of 16<sup>th</sup> to 1200 UTC of 18<sup>th</sup> May), AMPHAN moved slower than the average. Thereafter the speed increased and reached the maximum (29 kmph at 0600 UTC of 20<sup>th</sup> May) just prior to landfall.
- x. The Velocity Flux, Accumulated Cyclone Energy (a measure of damage potential) and Power Dissipation Index (a measure of loss) were  $16.30 \times 10^2$  knots,  $15.45 \times 10^4$  knots<sup>2</sup> and  $16.04 \times 10^6$  knots<sup>3</sup> respectively against the LPA during 1990-2013 of  $5.28 \times 10^2$  knots,  $8.6 \times 10^4$  knots<sup>2</sup> and  $2.8 \times 10^6$  knots<sup>3</sup> respectively for tropical cyclones over BoB during pre-monsoon season.



**Fig.1: Observed track of SuCS ‘AMPHAN’ over the southeast Bay of Bengal (16-21 May, 2020)**



**Fig.2: Tracks of severe cyclonic storms and above intensity storms (a) developing in the grid  $\pm 2.5^\circ$  of genesis point and (b) crossing West Bengal-Bangladesh coasts in the month of May during satellite era 1960 onwards**



**Fig. 3: (a) Translational speed & direction of movement and (b) Maximum sustained surface winds (kts) & Estimated Central Pressure**

The six hourly average translational speed during the life cycle of SuCS Amphan is presented in Fig. 3a. The six hourly maximum sustained wind speed and estimated central pressure is presented in Fig. 3b. After landfall, the system exhibited rapid weakening during 1500 UTC of 20<sup>th</sup> to 0600 UTC of 21<sup>st</sup> May.

**Table 1: Best track positions and other parameters of the Super Cyclonic Storm, ‘AMPHAN’ over the Bay of Bengal during 16 May- 21 May, 2020**

Date	Time (UTC)	Centre lat. <sup>0</sup> N/ long. <sup>0</sup> E	C.I. NO.	Estimated Central Pressure (hPa)	Estimated Maximum Sustained Surface Wind (kt)	Estimated Pressure drop at the Centre (hPa)	Grade
16/05/2020	0000	10.4   87.0	1.5	1000	25	03	D


	0300	10.7	86.5	1.5	1000	25	03	D	
	0600	10.9	86.3	1.5	1000	25	03	D	
	0900	10.9	86.3	2.0	998	30	05	<b>DD</b>	
	1200	10.9	86.3	2.5	996	35	07	<b>CS</b>	
	1500	11.0	86.2	2.5	995	40	08	CS	
	1800	11.1	86.1	2.5	995	40	08	CS	
	2100	11.3	86.1	3.0	994	45	10	CS	
17/05/2020	0000	11.4	86.0	3.0	992	45	10	CS	
	0300	11.4	86.0	3.0	990	50	12	<b>SCS</b>	
	0600	11.5	86.0	3.5	988	55	15	SCS	
	0900	11.7	86.0	4.0	980	65	22	<b>VSCS</b>	
	1200	12.0	86.0	4.0	978	70	25	VSCS	
	1500	12.8	86.2	4.0	978	70	25	VSCS	
	1800	12.5	86.1	4.5	970	80	32	VSCS	
2100	12.9	86.4	5.0	962	90	40	<b>ESCS</b>		
18/05/2020	0000	13.2	86.3	5.5	952	100	50	ESCS	
	0300	13.3	86.2	6.0	936	115	66	ESCS	
	0600	13.4	86.2	6.5	930	120	72	<b>SuCS</b>	
	0900	13.7	86.2	6.5	930	120	72	SuCS	
	1200	14.0	86.3	6.5	926	125	76	SuCS	
	1500	14.5	86.4	6.5	926	125	76	SuCS	
	1800	14.9	86.5	6.5	920	130	84	SuCS	
2100	15.2	86.6	6.5	920	130	84	SuCS		
19/05/2020	0000	15.6	86.7	6.5	926	125	76	SuCS	
	0300	16.0	86.8	6.0	930	120	72	SuCS	
	0600	16.5	86.9	6.0	936	115	66	<b>ESCS</b>	
	0900	17.0	86.9	6.0	942	110	60	ESCS	
	1200	17.4	87.0	5.5	946	105	56	ESCS	
	1500	18.1	87.1	5.5	950	100	50	ESCS	
	1800	18.4	87.2	5.5	950	100	50	ESCS	
2100	18.7	87.2	5.5	950	100	50	ESCS		
20/05/2020	0000	19.1	87.5	5.0	954	95	46	ESCS	
	0300	19.8	87.7	5.0	958	90	40	ESCS	
	0600	20.6	88.0	5.0	960	90	40	ESCS	
	0900	21.4	88.1	5.0	960	90	40	ESCS	
	Crossed West Bengal – Bangladesh coasts as a very severe cyclonic storm across Sundarbans, near lat.21.65°N/long. 88.3°E during1000-1200 UTC, with maximum sustained wind speed of 85 knots gusting to 100 knots.								
	1200	21.9	88.4	-	968	80	32	<b>VSCS</b>	
	1500	22.7	88.6	-	978	65	32	VSCS	
	1800	23.3	89.0	-	984	50	14	<b>SCS</b>	
	2100	24.2	89.0	-	986	45	12	<b>CS</b>	
21/05/2020	0000	24.2	89.3	-	988	40	10	CS	
	0300	24.7	89.5	-	990	35	08	CS	
	0600	25.0	89.6	-	992	30	06	<b>DD</b>	

	1200	25.4	89.6	-	995	20	04	<b>D</b>
	1800	Weakened into a well marked low pressure area over north Bangladesh and neighbourhood						

### 3. Monitoring and Prediction:

India Meteorological Department (IMD) maintained round the clock watch over the north Indian Ocean and the system was monitored since 23<sup>rd</sup> April about three weeks prior to the formation of the Low Pressure Area over the southeast BoB on 13<sup>th</sup> May. A Low Pressure Area formed over south Andaman Sea and adjoining southeast Bay of Bengal on 1<sup>st</sup> May. It meandered over the region for next 5 days and became less marked on 6<sup>th</sup>. However, associated cyclonic circulation persisted over the region till 12<sup>th</sup>. In the extended range outlook issued on 7<sup>th</sup> May, IMD indicated possible Cyclogenesis during the second week over south Andaman Sea and adjoining southeast Bay of Bengal. Accordingly, continuous watch of this circulation was maintained. On 9<sup>th</sup> May, it was indicated that a Low Pressure Area would form over the region on 13<sup>th</sup> May (96 hours prior to formation of the system) under the influence of the remnant cyclonic circulation persisting over the region during 6<sup>th</sup>-12<sup>th</sup>. On 11<sup>th</sup>, it was indicated that cyclogenesis (formation of depression) would occur around 16<sup>th</sup> May (48 hours prior to formation of the Low Pressure Area and 120 hours prior to formation of depression) over the BoB.

The cyclone was monitored with the help of available satellite observations from INSAT 3D and 3DR, polar orbiting satellites including SCATSAT, ASCAT etc. and available ships & buoy observations in the region. From 18<sup>th</sup> May midnight (1800 UTC) onwards till 20<sup>th</sup> May, the system was tracked gradually by IMD Doppler Weather Radars (DWRs) at Visakhapatnam, Gopalpur, Paradip, Kolkata and Agartala as it moved from south to north. IMD also utilised DWR products from 'DRDO Integrated Test Range', Chandipur, Balasore for tracking the system. Various numerical weather prediction models run by Ministry of Earth Sciences (MoES) institutions, various global models and IMD's dynamical-statistical models developed in-house were utilized to predict the genesis, track, landfall and intensity of the cyclone. A digitized forecasting system of IMD was utilized for analysis and comparison of guidance from various models, decision making process and warning product generation. Typical satellite and radar imageries are presented in **Fig. 4**. The satellite and radar imageries during entire life cycle of the system are placed at Annexure-1.




**Fig.4: Typical satellite imageries for SuCS Amphan over the Bay of Bengal and Radar imageries from DWR Kolkata & Paradip**

#### **4. Realized Weather:**

##### **4.1. Realised rainfall**

Rainfall associated with SuCS AMPHAN based on IMD-NCMRWF GPM merged gauge 24 hours cumulative rainfall ending at 0830 IST of date is depicted in **Fig 5**.

It indicates occurrence of heavy to very heavy rainfall at a few places over coastal Odisha & Gangetic West Bengal on 20<sup>th</sup> May, heavy rainfall at isolated places over Gangetic West Bengal & adjoining Bangladesh and Assam, Meghalaya & Arunachal Pradesh on 21<sup>st</sup> May and heavy rainfall at isolated places over Assam, Meghalaya, Arunachal Pradesh, Sikkim, Nagaland, Manipur & Mizoram on 22<sup>nd</sup> May.



**Fig.5: IMD-NCMRWF GPM merged gauge 24 hr cumulative rainfall (cm) ending at 0830 IST of date during 16<sup>th</sup> May – 22<sup>nd</sup> May and 7 days average rainfall (cm/day)**

Realized 24 hrs accumulated rainfall ( $\geq 7$ cm) ending at 0830 hrs IST of date during the life cycle of the system is presented below:

### 20 May 2020

**Gangetic West Bengal:** Contai-11, Digha-9,

**Odisha:** Paradip-21, Balikuda-18, Kakatpur & Kujanga-16 each, Astaranga & Alipingal-14 each, Niali-12, Raghunathpur, Puri, Marsaghai, Nilgiri, Kantapada & Garadapur-9 each, Gop, Chandanpur, Betanati, Rajkanika, Jagatsinghpur, Tirtol & Baripada-8 each and Binjharpur, Satyabadi, Nischintakoili, Chandbali, Bhograi, Jajpur, Dhamnagar, Soro, Tihidi, Bari & Basudevpur-7 each,

### 21 May 2020

**Assam & Meghalaya:** Williamnagar-23, Mawsynram-15, Sohra (RKM)-13, Bhaghmara-11 and Sohra & Shillong-9 each

**SHWB & Sikkim:** Sevoke-7,

**Gangetic West Bengal:** Alipore-24, Dum Dum-20, Harinkhola & Debagram-13 each, Burdwan-10, Manteswar & Digha-9 each and Mohanpur, Kharagpur, Suri, Mangalkote, Bankura, Lalgah & Midnapore-7 each,

**Odisha:** Bhograi-13, Rajghat & Jaleswar-12 each and Chandanpur & Bangiriposi-11 each, Paradip, Samakhunta, Betanati & Baripada-9 each and Chandikhol, Joshipur & Danagadi-7 each

22 May 2020

**Arunachal Pradesh:** Bhalukpong-12, Bomdila-10, Itanagar & Ziro-9 each, Roing-8, Pasighat Aero-7,

**Assam & Meghalaya:** Cherrapunji (RKM)-25, Mawsynram-22, Sohra -21, Khliehriat-15, Goalpara-12, Nongstein, Dhubri & Mela bazar -10 each, Goibargaon, N.Lakhimpur/Lilabari & Shillong-8 each and Tamulpur, Barapani, Williamnagar, DRF & Tezpur-7 each,

**Nagaland, Manipur, Mizoram & Tripura:** Sabroom-13,


**SHWB & Sikkim:** Buxaduar-8,

#### 4.2. Realised wind:

Kolkata (Dum Dum) reported 130 kmph at 1855 hrs IST (1325 UTC) and Kolkata (Alipore) 112 kmph at 1752 hrs IST (1222 UTC) of 20th May.


Also Paradip reported 106 kmph at 0630 hrs IST ( 0100 UTC) , Chandbali, 80 kmph at 0830 hrs.IST (0300 UTC) and Balasore 91 kmph during 1330 – 1430 hrs. IST (0800 & 0900 UTC) of 20<sup>th</sup> May.

Maximum sustained wind speed recordings from Automated Weather Stations (AWS) at Canning, Nimpith and Sagar Islands on 20<sup>th</sup> May are shown in **Fig.6**.



**Fig.6:** AWS data from Canning, Sagar Island & Nimpith on 20<sup>th</sup> May 2020

Five hours period during 1600-2100 hrs IST witnessed wind speed 60-80 kmph with many gusts of 100 kmph (with maximum of 114 kmph) over Kolkata. **Fig.7** indicates the wind recorded at Alipore (Kolkata) on 20th May 2020.



**Fig.7:** Wind anemograph recordings at Kolkata (Alipore) on 20<sup>th</sup> May 2020

#### **4.3. Realised storm surge:**

As per the Post Cyclone landfall survey conducted by Area Cyclone Warning Centre (ACWC) Kolkata, Tidal waves of 15 Feet height inundated low lying areas of South & North 24 Parganas and adjoining areas of east Medinipur Districts of West Bengal.

#### **5. Forecast performance:**

##### **5.1. Genesis Forecast**


- The system was monitored since 23<sup>rd</sup> April about three weeks prior to the formation of Low Pressure Area over the southeast BoB on 13<sup>th</sup> May. A Low Pressure Area formed over south Andaman Sea and adjoining southeast Bay of Bengal on 1<sup>st</sup> May. It meandered over the region for next 5 days and became less marked on 6<sup>th</sup>. However, associated cyclonic circulation persisted over the region till 12<sup>th</sup>. Continuous watch of this circulation was maintained.
- In the extended range outlook issued on 7<sup>th</sup> May, cyclogenesis (formation of Depression) was predicted with low probability in the later part of week during 8<sup>th</sup>-14<sup>th</sup> May 2020. It was also predicted that the system would intensify further and move initially north-northwestwards and recurve north-northeastwards thereafter towards north BoB.
- In the Tropical Weather Outlook issued on 9<sup>th</sup> May, it was indicated that a Low Pressure Area would form over the region on 13<sup>th</sup> May (96 hours prior to formation of the system) under the influence of the remnant cyclonic circulation persisting over the region during 6<sup>th</sup>-12<sup>th</sup>.

- In the Tropical Weather Outlook issued on 11<sup>th</sup> May, it was indicated that cyclogenesis (formation of depression) would occur around 16<sup>th</sup> May (48 hours prior to formation of the Low Pressure Area and 120 hours prior to formation of depression) over the BoB. The Low pressure area formed on 13<sup>th</sup> May and concentrated into a Depression on 16<sup>th</sup> May morning.

## 5.2. Track, Intensity and Landfall Forecast

- First information was provided in the **extended range outlook issued on 7th May (about 6 days prior to formation of LPA, 9 days prior to formation of depression and 13 days prior to Landfall)** indicated that the system would intensify into a cyclonic storm and move initially northwestwards and recurve north-northeastwards towards north BoB
- In the **Tropical Weather Outlook, Press release and inforatory message to the Government of India issued on 13th April (on the day of development of LPA, 3 days prior to formation of depression and 7 days prior to Landfall)**, it was indicated that the system would intensify into a cyclonic storm by 16th evening and would move initially northwestwards till 17th and then recurve north-northeastwards towards north BoB.
- Actually, the depression formed in the morning (0000 UTC) of 16th, cyclonic storm in the evening (1200 UTC) of 16th and the system moved north-northwestwards till 17th evening (1200 UTC) followed by north-northeastward recurvature thereafter and crossed West Bengal coast on 20th Afternoon.
- **In the first bulletin issued at 0845 IST of 16<sup>th</sup> May (104 hrs prior to landfall)** with the formation of Depression, it was indicated that the system would intensify into a cyclonic storm and will move north-northwestwards till 17<sup>th</sup> May followed by north-northeastward re-curved towards West Bengal coast during 18<sup>th</sup>-20<sup>th</sup> May and cross West Bengal coast with maximum sustained wind speed of 155-165 kmph gusting to 180 kmph.
- **In the bulletin issued at 1645 hrs IST of 16<sup>th</sup> May (24 hrs prior to rapid intensification)**, rapid intensification of the system was predicted and the system rapidly intensified from 17<sup>th</sup> afternoon onwards.
- **In the bulletin issued at 0845 hrs IST of 17<sup>th</sup> May (80 hrs prior to landfall)**, it was precisely mentioned that the system would cross West Bengal-Bangladesh coasts between Sagar Island (West Bengal) and Hatiya Islands (Bangladesh coast) during afternoon to evening of 20<sup>th</sup> May with maximum sustained wind speed of 155-165 kmph gusting to 185 kmph. The predicted track indicated Landfall across Sunderbans on 20<sup>th</sup> Afternoon.
- Observed & forecast track based on 0600 UTC of 17<sup>th</sup> May, 0300 UTC of 18<sup>th</sup> May and 0300 UTC of 19<sup>th</sup> May about 84, 60 & 36 hrs prior to landfall respectively of SuCS Amphan indicating accurate track, landfall and intensity prediction is presented in **Fig.8 (a-c)**. It also indicates the consistency in the forecast track & intensity of the Cyclone.

- IMD continuously predicted since 16<sup>th</sup> May that Amphan will cross West Bengal coast as a very severe cyclonic storm (VSCS) with wind speed of 155-165 kmph gusting to 180 kmph on 20<sup>th</sup> May.



**Fig.8(a): Observed and forecast track along with cone of uncertainty and quadrant wind distribution based on 0600 UTC of 17<sup>th</sup> May (84 hrs prior to landfall) of SuCS AMPHAN indicating accuracy in landfall, track & intensity predictions**





Fig.8(b) &(c): Observed and forecast track alongwith cone of uncertainty and quadrant wind distribution based on (b) 0300 UTC of 18<sup>th</sup> May (60 hours prior to landfall) and (c) 0300 UTC of 19<sup>th</sup> May (36 hours prior to landfall) of SuCS AMPHAN indicating accuracy in landfall, track & intensity predictions

### 5.3. Landfall Forecast Errors:

The landfall point and time Forecast errors compared to long period average (LPA) errors during 2015-19 are presented in Table 2 and Fig. 9 (a-b).

- The landfall point forecast errors for 24, 48 and 72 hrs lead period were 5.5, 11.0, and 35.2 km respectively against the LPA errors of 44.7, 69.4 and 109.3 km during 2015-19 respectively (**Fig. 9a**).
- The landfall time forecast errors for 24, 48 and 72 hrs lead period were 0.5, 0, and 2.0 hours respectively against the LPA errors of 3.0, 5.4 and 8.6 hours during 2015-19 respectively (**Fig. 9b**).
- **For all lead periods, the landfall errors were exceptionally less than the LPA errors during 2015-19. Considering the Eye (Centre) Diameter of Cyclone ‘Amphan’ as 15 km at the time of landfall, there was almost zero error in landfall time and point forecast upto 48 hrs lead period.**



**Fig. 9: Landfall (a) point and (b) time forecast errors of SUCS AMPHAN as compared to long period average (2015-19)**

**Table 2: Landfall point and time forecast errors of SuCS Amphan as compared to long period average (LPA) errors during 2015-19**


Lead Period (hrs)	Base Time	Landfall Point ( <sup>0</sup> N/ <sup>0</sup> E)		Landfall Time (hours)		Operational Error		LPA error (2015-19)	
		Forecast	Actual	Forecast	Actual	LPE (km)	LTE (hours)	LPE (km)	LTE (hours)
12	20/00	21.65/88.30	21.65/88.30	20/1100	20/1100	0.0	0.0	25.4	2.0
24	19/12	21.70/88.30	21.65/88.30	20/1130	20/1100	5.5	+0.5	44.7	3.0
36	19/00	21.65/88.35	21.65/88.30	20/1130	20/1100	5.5	+0.5	41.7	4.7
48	18/12	21.65/88.40	21.65/88.30	20/1100	20/1100	11.0	0.0	69.4	5.4
60	18/00	21.65/88.40	21.65/88.30	20/0930	20/1100	11.0	-1.5	89.6	5.5
72	17/12	21.90/88.10	21.65/88.30	20/1300	20/1100	35.2	+2.0	109.3	8.6
84	17/00	21.63/88.87	21.65/88.30	20/1200	20/1100	62.7	+1.0	141.3	5.8
96	16/12	21.80/87.90	21.65/88.30	20/1000	20/1100	47.0	-1.0	210.5	8.8
108	16/00	21.80/87.90	21.65/88.30	20/1200	20/1100	47.0	+1.0	350.8	9.0

“+” indicates delayed prediction and “-” indicates early prediction

#### 5.4. Track Forecast Errors:

The track forecast errors (Forecast position – Actual position of Cyclone centre) and skill as compared to Climatological and Persistence forecast are presented in Table 3 and Fig. 10 (a-b).

- The track forecast errors for 24, 48 and 72 hrs lead period were 59.4, 59.9, and 61.0 km respectively against the LPA errors of 80.6, 125.5, and 171.2 km respectively (**Fig.10a**).
- The track forecast skill was about 60%, 78%, and 88% against the LPA skill of 61%, 73%, and 74% for 24, 48 and 72 hrs lead period respectively (**Fig.10b**).
- **Track forecast errors were exceptionally less than the past five years average errors for all lead periods. Similarly, track forecast skill was higher than the past five years average skill for all lead periods beyond 24 hours.**



**Fig. 10(a) & (b): Track forecast errors and skill of SUCS AMPHAN as compared to long period average (2015-19)**


**Table 3: Operational track forecast errors (km) & Skill (%) compared to long period average during 2015-19**

Lead Period (hrs)	No. of obs. Verified	Operational Track Forecast		Long Period Average (2015-19) Track Forecast	
		Error (km)	Skill (%)*	Error (km)	Skill (%)*
12	18	49.0	48.6	49.6	57.8
24	16	59.4	59.5	80.6	60.9
36	15	55.4	72.5	99.9	68.5
48	13	59.9	78.3	125.5	72.6
60	11	61.9	85.2	147.8	73.0
72	9	61.0	88.0	171.2	74.2
84	7	105.5	84.0	208.3	74.4
96	5	88.8	90.3	240.5	75.9
108	3	86.3	92.1	259.8	75.6
120	1	93.3	92.6	322.2	76.0

### 5.5. Intensity Forecast Errors:


The intensity forecast errors (Forecast wind – Actual wind) and skill based on absolute errors and root mean square errors are presented in Table 4 and Fig. 11 & 12 respectively.

- The absolute error (AE) of intensity (wind) forecast for 24, 48 and 72 hrs lead period were 13.4, 15.0 and 9.0 knots against the LPA errors of 8.9, 13.0, and 15.4 knots during 2015-19 respectively (**Fig. 11**).
- The root mean square error (RMSE) of intensity (wind) forecast for 24, 48 and 72 hrs lead period were 18.8, 22.1 and 12.2 knots against the LPA errors of 11.5, 16.7, and 19.2 knots respectively (**Fig. 11**).



**Fig. 11: Absolute errors (AE) and Root Mean Square errors (RMSE) in intensity forecast (winds in knots) of SUCS AMPHAN as compared to long period average (2015-19)**

- The skill (%) in intensity forecast as compared to persistence forecast based on AE for 24, 48 and 72 hrs lead period was 56%, 80% and 94% against the LPA of 45%, 69% and 72% respectively (**Fig. 12a**).
- The skill (%) in intensity forecast based on RMSE for 24, 48 and 72 hrs lead period was 50%, 76% and 93% against the LPA of 49%, 63% and 72% respectively (**Fig. 12b**).
- Thus even being a Super cyclone, the intensity forecast was skill full and better than the Long Period Average.



**Fig. 12: Skill (%) in intensity forecast based on (a) Absolute errors (AE) and (b) Root Mean Square errors (RMSE) of SUCS AMPHAN as compared to long period average (2015-19)**

**Table 4: Mean Intensity forecast errors (kt) and Skill (%) in association with SUCS AMPHAN compared to long period average errors and skill during 2015-19**

Lead Period (hrs)	N	Average error in Intensity forecast (kts)		LPA (2015-19) Intensity forecast error (kts)		Operational Skill* (%) in intensity forecast		LPA (2015-19) Skill* (%) in intensity forecast	
		AE	RMSE	AE	RMSE	AE	RMSE	AE	RMSE
12	18	7.7	10.5	5.4	7.1	43.1	38.6	28.3	36.3
24	16	13.4	18.8	8.9	11.5	55.8	50.4	44.9	49.3
36	15	16.9	23.8	12.2	14.9	66.2	61.0	64.6	55.5
48	13	15.0	22.1	13.0	16.7	79.7	76.0	68.5	62.9
60	11	11.9	17.0	14.2	17.6	89.1	87.6	70.3	69.6
72	9	9.0	12.2	15.4	19.2	93.5	93.2	72.0	72.1
84	7	10.8	12.5	16.7	20.4	91.8	92.4	74.8	76.4
96	5	13.9	15.6	17.2	20.5	88.8	89.2	77.7	80.2
108	3	8.7	10.1	16.6	20.0	92.4	91.4	81.9	84.2
120	1	4.3	4.3	16.4	20.1	95.7	95.7	83.8	85.3

N: No. of observations verified; AE: Absolute Error; RMSE: Root Mean Square Error, LPA: Long Period Average (2014-18).

### 5.6. Adverse weather forecast verification

The verifications of adverse weather like heavy rainfall, gale wind and storm surge forecast issued by IMD are presented in Tables 5-7. It is found that all the three types of adverse weather were predicted accurately and well in advance.

**Table 5: Verification of Heavy Rainfall Forecast**

Date/Base Time of forecast (UTC)	24 hr Heavy rainfall warning ending at 0300 UTC of next day	Realised 24-hour heavy rainfall ending at 0300 UTC of date
16.05.2020/0300	<p>Andaman &amp; Nicobar Islands</p> <p>Heavy falls at isolated places is very likely over Andaman &amp; Nicobar Islands on 16th May.</p> <p>Odisha &amp; Gangetic West Bengal</p> <p>Heavy falls at isolated places from 18th May evening, heavy to very heavy rainfalls at a few places on 19th May and isolated heavy rainfall over northeast Odisha on 20th May 2020. Coastal districts of Gangetic West Bengal are likely to experience heavy falls at a few places on 19th May, heavy to very heavy falls at a few places with extremely heavy falls at isolated places over Gangetic West Bengal on 20th May.</p>	<p>17 May 2020</p> <p><b>A &amp; N Island:</b></p> <p>Nancowary-2 and IAF Carnicobar &amp; Car Nicobar-1 each</p> <p>18 May 2020</p> <p><b>A &amp; N Island:</b></p> <p>Car Nicobar-5, IAF Carnicobar-4 and Nancowary-1,</p> <p><b>Odisha:</b> Jhumpura-10 and Gurundia -8</p> <p>19 May 2020</p>
17.05.2020/0300	<p>(i) Rainfall (over Andaman &amp; Nicobar Islands):</p> <ul style="list-style-type: none"> <li>•Light to moderate rainfall at most places is very likely over Andaman &amp; Nicobar Islands during next 24 hours.</li> <li>•Rainfall (over Odisha &amp; Gangetic West Bengal)</li> </ul> <p>Heavy falls at isolated places from 18th May evening. Rainfall at most places with heavy to very heavy rainfall at a few places on 19th May and isolated heavy rainfall over north coastal Odisha on 20th May 2020.</p> <p>Coastal districts of Gangetic West Bengal are likely to experience heavy falls at isolated places on 19th May.</p> <p>Heavy to very heavy falls at a few places &amp; extremely heavy falls at isolated places likely over coastal districts of West Bengal on 20th May.</p>	<p>SHWB &amp; Sikkim:</p> <p>Mangan-7</p> <p>20 May 2020</p> <p><b>Gangetic West Bengal:</b></p> <p>Contai-11, Digha-9,</p> <p><b>Odisha:</b> Paradeep-21, Balikuda-18, Kakatpur &amp; Kujanga-16 each, Astaranga &amp; Alipingal-14 each, Niali-12, Raghunathpur, Puri, Marsaghai, Nilgiri, Kantapada &amp; Garadapur-9 each, Gop, Chandanpur, Betanati, Rajkanika, Jagatsinghpur, Tirtol &amp; Baripada-8 each and Binjharpur, Satyabadi, Nischintakoili, Chandbali, Bhograi, Jajpur, Dhamnagar, Soro, Tihidi, Bari &amp; Basudevpur-7 each,</p> <p>21 May 2020</p> <p><b>Assam &amp; Meghalaya:</b></p> <p>Williamnagar-23, Mawsynram-15, Sohra</p>
18.05.2020/0300	<p><b>Odisha</b></p> <p>Coastal Odisha is likely to experience heavy falls at isolated places over coastal Odisha (Gajapati, Ganjam, Puri, Jagatsinghpur &amp; Kendrapara Districts) on 18th May, 2020.</p> <p>Heavy to very heavy rainfall at a few places over north coastal Odisha (Jagatsinghpur, Kendrapara, Jajpur, Balasore, Bhadrak &amp; Mayurbhanj Districts) and isolated heavy falls over Khordha &amp; Puri districts on 19th May and isolated heavy rainfall over north Odisha (Bhadrak, Balasore, Mayurbhanj, Jajpur, Kendrapara and Keonjhar Districts) on 20th May 2020.</p> <p><b>West Bengal</b></p> <p>Coastal districts of Gangetic West Bengal (East Medinipur, South &amp; North 24 Parganas) are likely to experience heavy falls at isolated places on</p>	<p>Chandanpur, Betanati, Rajkanika, Jagatsinghpur, Tirtol &amp; Baripada-8 each and Binjharpur, Satyabadi, Nischintakoili, Chandbali, Bhograi, Jajpur, Dhamnagar, Soro, Tihidi, Bari &amp; Basudevpur-7 each,</p> <p>21 May 2020</p> <p><b>Assam &amp; Meghalaya:</b></p> <p>Williamnagar-23, Mawsynram-15, Sohra</p>

	<p>19th May. Rainfall at most places with heavy to very heavy falls at a few places &amp; extremely heavy falls at isolated places likely over Gangetic West Bengal (east &amp; west Medinipur, south &amp; north 24 Parganas, Howrah, Hoogli, Kolkata and adjoining districts) on 20th May and isolated heavy rain over interior districts on 21st May, 2020.</p> <p><b>Sub-Himalayan West Bengal and Sikkim</b> Heavy to very heavy falls at a few places over Malda &amp; Dinajpur districts on 20th May and over most of the districts of Sub-Himalayan West Bengal &amp; Sikkim on 21st May, 2020.</p> <p><b>Assam &amp; Meghalaya</b> Heavy to very heavy falls at a few places over the western districts of Assam &amp; Meghalaya on 21st May.</p>	<p>(RKM)-13, Bhaghmara-11 and Sohra &amp; Shillong-9 each</p> <p><b>SHWB &amp; Sikkim:</b> Sevoke-7, <b>Gangetic West Bengal:</b> Alipore-24, Dum Dum-20, Harinkhola &amp; Debagram-13 each, Burdwan-10, Manteswar &amp; Digha-9 each and Mohanpur, Kharagpur, Suri, Mangalkote, Bankura, Lalgarh &amp; Midnapore-7 each,</p> <p><b>Odisha:</b> Bhograi-13, Rajghat &amp; Jaleswar-12 each and Chandanpur &amp; Bangiriposi-11 each, Paradeep, Samakhunta, Betanati &amp; Baripada-9 each and Chandikhol, Joshipur &amp; Danagadi-7 each,</p> <p>22 May 2020</p> <p><b>Arunachal Pradesh:</b> Bhalukpong-12, Bomdila-10, Itanagar &amp; Ziro-9 each, Roing-8, Pasighat Aero-7,</p> <p><b>Assam &amp; Meghalaya:</b> Sohra (RKM)-25, Mawsynram-22, Sohra-21, Khliehriat-15, Goalpara-12, Goalpara-11, Nongstein, Dhubri &amp; Mela bazar -10 each, Goibargaon, N.Lakhimpur/Lilabari &amp; Shillong-8 each and Tamulpur, Barapani, Williamnagar, DRF &amp; Tezpur-7 each,</p> <p><b>Nagaland, Manipur, Mizoram &amp; Tripura:</b></p>
19.05.2020/0300	<p><b>Odisha</b> Coastal Odisha to experience heavy to very heavy rainfall at isolated places Jagatsinghpur, Kendrapara and Bhadrak Districts and isolated heavy falls over Jajpur, Balasore, Cuttack, Mayurbhanj, Khordha &amp; Puri districts on 19th May and isolated heavy to very heavy rainfall at isolated places over north coastal Odisha (Jagatsinghpur, Bhadrak and Keonjhar Districts) on 20th May 2020.</p> <p><b>West Bengal</b> Coastal districts of Gangetic West Bengal (East Medinipur, South &amp; North 24 Parganas) are likely to experience light to moderate rainfall at many places commencing from today, the 19th May afternoon. Rainfall intensity is likely to increase gradually and become maximum on 20th May. Heavy to very heavy falls at a few places &amp; extremely heavy falls at isolated places likely over Gangetic West Bengal (east &amp; west Medinipur, south &amp; north 24 Parganas, Howrah, Hoogli, Kolkata and adjoining districts) on 20th May and isolated heavy rain over interior districts on 21st May, 2020.</p> <p><b>Sub-Himalayan West Bengal and Sikkim</b> Heavy to very heavy falls at a few places over Malda &amp; Dinajpur districts on 20th May and over most of the districts of Sub-Himalayan West Bengal &amp; Sikkim on 21st May, 2020.</p> <p><b>Assam &amp; Meghalaya</b> Heavy to very heavy falls at a few places over the western districts of Assam &amp; Meghalaya on 21st May.</p>	
20.05.2020/0300	<p><b>Odisha</b> Heavy to very heavy rainfall very likely at isolated places over north coastal Odisha (Balasore,</p>	

	<p>Bhadrak, Mayurbhanj, Jajpur, Kendrapara and Keonjhar Districts) and isolated heavy falls over Jagatsinghpur district on 20th May 2020.</p> <p><b>West Bengal</b> Heavy to very heavy falls at a few places &amp; extremely heavy falls at isolated places likely over Gangetic West Bengal (east &amp; west Medinipur, south &amp; north 24 Parganas, Howrah, Hoogli, Kolkata and adjoining districts) on 20th May and isolated heavy rain over interior districts on 21st May, 2020.</p> <p><b>Sub-Himalayan West Bengal and Sikkim</b> Heavy to very heavy falls at a few places over Malda &amp; Dinajpur districts on 20th May and over most of the districts of Sub-Himalayan West Bengal &amp; Sikkim on 21st May, 2020.</p> <p><b>Assam &amp; Meghalaya</b> Heavy to very heavy falls at a few places over the western districts of Assam &amp; Meghalaya on 21st May.</p>	<p>Sabroom-13, <b>SHWB &amp; Sikkim:</b> Buxaduar-8,</p>
21.05.2020/0300	<p><b>Assam &amp; Meghalaya</b> Heavy to very heavy &amp; extremely heavy falls (<math>\geq 20</math> cm) at isolated places very likely on 21st May, 2020.</p> <p><b>Arunachal Pradesh</b> Heavy to very heavy falls at isolated places very likely on 21st May, 2020.</p>	

**Table 6: Verification of Squally/Gale wind forecast (16-21 May)**

Date/Base Time of Forecast (UTC)	Gale/ Squally wind Forecast at 0300 UTC of date	Realised wind
16.05.2020/0300	<ul style="list-style-type: none"> <li>Squally wind speed reaching 45 to 55 kmph gusting to 65 kmph is likely along and off Odisha coast from 19th morning and along and off West Bengal coast from 19th afternoon. The wind speed will gradually increase becoming gale wind speed reaching 75 to 85 kmph gusting to 95 kmph from 20th morning along and off north Odisha and West Bengal coast. It will gradually increase thereafter.</li> <li>Squally wind speed reaching 45 to 55 kmph gusting to 65 kmph is likely over Andaman Sea during next 48 hours.</li> <li>Squally wind speed reaching 45 to 55 kmph gusting to 65 kmph is prevailing over southeast and adjoining southwest Bay of Bengal. It is likely to increase becoming 90-100 gusting to 110 kmph over eastcentral and adjoining west central Bay of Bengal by 17th morning, 120-130 gusting to 145 kmph over southern parts of central Bay of Bengal by 18th morning, 155-165</li> </ul>	<p>155-165kmph gusting to 185 kmph along &amp; off coastal Districts of West Bengal and 100-110 kmph gusting to 120 kmph along &amp; off north coastal Districts of Odisha. Kolkata (Dum Dum) reported 130 kmph at 1855 hrs IST and Kolkata (Alipore) 112 kmph at 1752 hrs IST of 20th May. Also Canning reported 85 knots</p>

	gusting to 180 kmph over northern parts of central Bay of Bengal and adjoining north Bay of Bengal on 19th , and 160-170 gusting to 190 kmph over north Bay of Bengal by 20th morning.	9157 kmph), Nimpith 84 knots (155 kmph) and Sagar islands 60 knots (111 kmph)
17.05.2020/0300	<ul style="list-style-type: none"> <li>• Squally wind speed reaching 45 to 55 kmph gusting to 65 kmph is likely to commence along and off south Odisha coast from 18th evening, extend to along &amp; off north Odisha coast from 19th morning and along and off West Bengal coast from 19th afternoon. The wind speed will gradually increase becoming gale wind speed reaching 75 to 85 kmph gusting to 95 kmph from 20th morning along and off north Odisha and West Bengal coasts. It will gradually increase thereafter along &amp; off West Bengal coast.</li> <li>• Squally wind speed reaching 45 to 55 kmph gusting to 65 kmph is likely over Andaman Sea during next 24 hours.</li> <li>• Squally wind speed reaching 80 to 90 kmph gusting to 100 kmph is prevailing over southeast and adjoining southwest Bay of Bengal. It is likely to increase becoming 125-135 gusting to 150 kmph over southern parts of central Bay of Bengal by 18th morning, 160-170 gusting to 190 kmph over northern parts of central Bay of Bengal and adjoining North Bay of Bengal on 19th, and 155-165 gusting to 180 kmph over north Bay of Bengal during 20th May.</li> </ul>	Paradip reported 106 kmph at 0630 hrs IST (0100 UTC) , Chandbali, 80 kmph at 0830 hrs. IST (0300 UTC) and Balasore 91 kmph during 1330 – 1430 hrs. IST ( 0800 & 0900 UTC) of 20th May.
18.05.2020/0300	<p>West Bengal &amp; Odisha</p> <ul style="list-style-type: none"> <li>• Squally wind speed reaching 45 to 55 kmph gusting to 65 kmph is likely to commence along and off south Odisha coast from 18th evening, increase becoming 55 to 65 kmph gusting to 75 kmph extend to along &amp; off north Odisha coast from 19th morning and along and off West Bengal coast from 19th afternoon.</li> <li>• The wind speed will gradually increase becoming gale wind speed reaching 75 to 85 kmph gusting to 95 kmph from 20th morning along and off north Odisha (Jagatsinghpur, Kendrapara, Bhadrak, Balasore and Mayurbhanj districts) and West Bengal (east &amp; west Medinipur, south &amp; north 24 Parganas, Howrah, Hoogli, Kolkata Districts). It will gradually increase thereafter becoming 110 to 120 kmph gusting to 135 kmph along &amp; off the above-mentioned districts of North Odisha.</li> <li>• Gale wind speed reaching 155 to 165 kmph gusting to 185 kmph very likely along &amp; off east Medinipur and north &amp; south 24 Parganas districts and 100-110 kmph gusting to 120 kmph over Kolkata, Hoogli, Howrah and West Mednipur Districts of West Bengal during the</li> </ul>	

	<p>time of landfall (20th afternoon to night).</p> <ul style="list-style-type: none"> <li>• Squally wind speed reaching 55-65 kmph gusting to 75 kmph likely to prevail over Puri, Khordha, Cuttack, Jajpur districts of Odisha during 20th May 2020.</li> </ul> <p>Deep Sea area</p> <ul style="list-style-type: none"> <li>• Gale wind speed reaching 210-220 gusting to 240 kmph is prevailing over west-central and adjoining central parts of south Bay of Bengal. It is likely to increase becoming 220-230 gusting to 250 kmph over northern parts of central Bay of Bengal and adjoining North Bay of Bengal from tonight to 19th May morning, likely to increase further becoming 230-240 gusting to 255 kmph by tonight.</li> <li>• Gale wind speed reaching 220-230 gusting to 255 kmph over north Bay of Bengal from 19th morning and gradually decrease becoming 155-165 kmph gusting to 180 kmph by 20th evening.</li> </ul>	
19.05.2020/0300	<p>West Bengal &amp; Odisha</p> <ul style="list-style-type: none"> <li>• Squally wind speed reaching 45 to 55 kmph gusting to 65 kmph is prevailing along &amp; off south Odisha coast. It is very likely to increase and extend northwards becoming 55 to 65 kmph gusting to 75 kmph northwards to north Odisha coast by today afternoon and along &amp; off West Bengal coast by tonight.</li> <li>• The wind speed will gradually increase becoming gale wind speed reaching 75 to 85 kmph gusting to 95 kmph from 20th morning along and off north Odisha (Jagatsinghpur, Kendrapara, Bhadrak, Balasore and Mayurbhanj districts) coast and West Bengal (east &amp; west Medinipur, south &amp; north 24 Parganas, Howrah, Hoogli, Kolkata Districts) coast. It will gradually increase thereafter becoming 100 to 110 kmph gusting to 125 kmph along &amp; off the above mentioned districts of North Odisha.</li> <li>• Gale wind speed reaching 155 to 165 kmph gusting to 185 kmph very likely along &amp; off east Medinipur and north &amp; south 24 Parganas districts and 110-120 kmph gusting to 130 kmph over Kolkata, Hoogli, Howrah and West Medinipur Districts of West Bengal during the time of landfall (20th afternoon to night).</li> <li>• Squally wind speed reaching 55-65 kmph gusting to 75 kmph likely to prevail over Puri, Khordha, Cuttack, Jajpur districts of Odisha during 20th May 2020.</li> </ul> <p>Deep Sea area</p> <ul style="list-style-type: none"> <li>• Gale wind speed reaching 225-235 gusting to 255 kmph is prevailing over westcentral Bay of Bengal. It is likely to prevail over northern parts</li> </ul>	

	<p>of central Bay of Bengal and adjoining North Bay of Bengal during 19th May.</p> <ul style="list-style-type: none"> <li>• Gale wind speed reaching 180-190 gusting 210 kmph over north Bay of Bengal from 19th afternoon. It will gradually decrease becoming 165-175 kmph gusting to 195 kmph by 20th afternoon.</li> </ul>	
20.05.2020/0300	<p>West Bengal &amp; Odisha</p> <ul style="list-style-type: none"> <li>• Gale wind, speed reaching 90 to 100 kmph gusting to 110 kmph is prevailing along and off Jagatsinghpur, Kendrapara, Bhadrak, Balasore and Mayurbhanj districts of Odisha and squally wind speed reaching 55 to 65 kmph gusting to 75 kmph is prevailing along &amp; off remaining coastal districts of Odisha (Puri, Khordha, Cuttack, Jajpur) during 20th May 2020.</li> <li>• It will gradually increase becoming 100 to 110 kmph gusting to 125 kmph along &amp; off the above mentioned districts of North Odisha during forenoon to afternoon of 20th May 2020.</li> <li>• Gale wind speed reaching 80 to 90 kmph gusting to 100 kmph is prevailing along and off West Bengal coast (east &amp; west Medinipur, south &amp; north 24 Parganas, Howrah, Hoogli, Kolkata Districts).</li> <li>• Gale wind speed reaching 155 to 165 kmph gusting to 185 kmph very likely along &amp; off east Medinipur and north &amp; south 24 Parganas districts and 110-120 kmph gusting to 130 kmph over Kolkata, Hoogli, Howrah and West Medinipur Districts of West Bengal during the time of landfall (20th afternoon to night).</li> </ul> <p>Deep Sea area</p> <ul style="list-style-type: none"> <li>• Gale wind speed reaching 165-175 gusting to 195 kmph is prevailing over northwest and adjoining westcentral Bay of Bengal. Gale wind, speed reaching 150-160 gusting to 175 kmph is prevailing over northern parts of central Bay of Bengal. It will gradually decrease becoming 155-165 kmph gusting to 185 kmph over North Bay of Bengal during the evening hours of today, the 20th May.</li> </ul>	
21.05.2020/0300	<p>Squally wind speed reaching 50 to 60 kmph gusting to 70 kmph very likely over Western Assam &amp; Western Meghalaya till evening and reduce gradually thereafter.</p>	

**Table 7: Verification of Storm Surge Forecast**

Date/Base Time of observation	Storm Surge Forecast at 0300 UTC of date	Realised surge
18.05.2020/0300	Storm Surge of about 4-5 meters above Astronomical Tide is likely to inundate low lying areas of south & north 24 Parganas and about 3-4 meters over the low lying areas of East Medinipur District of West Bengal during the time of Landfall.	Maximum tidal wave of 4.6 meters height inundated the low lying areas of South & North 24 Parganas districts and adjoining areas of east Medinipur district of West Bengal as estimated by the post Cyclone landfall survey Team of ACWC Kolkata. No significant Storm Surge has been reported along Odisha coast.
19.05.2020/0300	Storm Surge of about 4-5 meters above Astronomical Tide is likely to inundate low lying areas of south & north 24 Parganas and about 3-4 meters over the low lying areas of East Medinipur District of West Bengal during the time of Landfall.	
20.05.2020/0300	Storm Surge of about 4-5 meters above Astronomical Tide is likely to inundate low lying areas of south & north 24 Parganas and about 3-4 meters over the low lying areas of East Medinipur District of West Bengal during the time of Landfall.	

## 6. Warning Services

### Bulletins issued by Cyclone Warning Division, New Delhi

- **Track, intensity and landfall forecast:** IMD continuously monitored, predicted and issued bulletins containing track, intensity, and landfall forecast for +06, +12, +18, +24, +36 and +48... +120 hrs lead period commencing from 16<sup>th</sup> May morning till the system weakened into a low pressure area. The above forecasts were issued from the stage of depression onwards along with the cone of uncertainty in the track forecast five times a day and every three hours during the cyclone period. The hourly updates were also provided 24 hours prior to landfall till the system maintained the intensity of cyclonic storm over West Bengal.
- **Cyclone structure forecast for shipping and coastal hazard management:** The radius of maximum wind and radii of MSW  $\geq 28$ ,  $\geq 34$ ,  $\geq 50$

and  $\geq 64$  knots wind in four quadrants of cyclone was issued every six hourly, commencing from 16<sup>th</sup> May morning giving forecast for +06, +12, +18, +24, +36 and +120 hrs lead period.

- **Four stage Warning:**

- In the first bulletin released at 0845 hrs IST of 16<sup>th</sup> May (**104 hrs prior to landfall**), **Pre-cyclone Watch** for West Bengal-north Odisha coasts was issued.
- The warnings were further upgraded and **Cyclone Watch** for West Bengal and north Odisha coasts was issued at 2030 hrs IST of 16<sup>th</sup> May (**92 hrs prior to landfall**).
- **Cyclone Alert (Yellow Message)** for West Bengal and north Odisha coasts was issued at 0840 hrs IST of 17<sup>th</sup> May (**80 hrs prior to landfall**).
- **Cyclone Warning (Orange Message)** for West Bengal and north Odisha coasts was issued at 0845 hrs IST of 18<sup>th</sup> May (**56 hrs prior to landfall**).
- **Post landfall outlook (Red Message)** for interior districts of Gangetic West Bengal, Assam and Meghalaya was issued at 2330 hrs IST of 19<sup>th</sup> May (**17 hrs prior to landfall**).

- **Adverse weather warning bulletins:** Adverse weather warning bulletins: The tropical cyclone forecasts alongwith expected adverse weather like heavy rain, gale wind and storm surge was issued with every three hourly update to central, state and district level disaster management agencies including MHA NDRF, NDMA for all concerned states along the east coast of India including Andaman & Nicobar Islands, Tamil Nadu, Andhra Pradesh, Odisha, West Bengal, Assam & Meghalaya, Manipur, Mizoram & Tripura. The bulletins also contained the suggested action for disaster managers and general public in particular for fishermen. These bulletins were also issued to Defence including Indian Navy & Indian Air Force.

- **Warning graphics:** The graphical display of the observed and forecast track with cone of uncertainty and the wind forecast for different quadrants were disseminated by email and uploaded in the RSMC, New Delhi website (<http://rsmcnewdelhi.imd.gov.in/>) regularly. The adverse weather warnings related to heavy rain, gale/squally wind & storm surge were also presented in graphics alongwith colour codes in the website.

- **Warning and advisory through social media:** Daily updates (every three hourly or whenever there was any significant change in intensity/track/landfall) were uploaded on Facebook and Twitter during the life period of the system since the development of low pressure area over BoB. However, from 20<sup>th</sup> morning (0000 UTC) onwards, hourly updates were issued and sent to disaster managers by email, uploaded on websites,

posted on Facebook and Twitter till the system maintained the intensity of cyclonic storm.

- **Press Conference, Press release and Media briefing:** Press and electronic media were given daily updates since inception of system through press release, e-mail, website and SMS. Joint Press conference was held by Director General of Meteorology (IMD), Director General (NDRF) on 19<sup>th</sup>, 20<sup>th</sup> & 21<sup>st</sup> May at National Media Centre. First press release was issued on 13<sup>th</sup> May, when the system was a low pressure area.
- **Warning and advisory for marine community:** The three/six hourly Global Maritime Distress Safety System (GMDSS) bulletins were issued by the Marine Weather Services division at New Delhi and bulletins for maritime interest were issued by Area cyclone warning centres of IMD at Chennai, Kolkata and Cyclone warning centres at Bhubaneswar and Visakhapatnam to ports, fishermen, coastal and high sea shipping community.
- **Fishermen Warning:** Regular warnings for fishermen for deep Sea BoB and the states of West Bengal, Odisha, Andhra Pradesh, Tamil Nadu, Kerala, Andaman & Nicobar Islands were issued since 13<sup>th</sup> May on the development of low pressure area over southeast BoB and adjoining south Andaman Islands.
- **Advisory for international Civil Aviation:** The Tropical Cyclone Advisory Centre (TCAC) bulletin for International Civil Aviation were issued every six hourly to all meteorological watch offices in Asia Pacific region for issue of significant meteorological information (SIGMET). It was also sent to Aviation Disaster Risk Reduction (ADRR) centre of WMO at Hong Kong.
- **Diagnostic and prognostic features of cyclone:** The prognostics and diagnostics of the systems were described in the RSMC bulletins.
- **Hourly Bulletin:** Hourly updates on the location, distance from recognised station, intensity and landfall commenced from 20<sup>th</sup> morning (0000 UTC) onwards till the system maintained the intensity of cyclonic storm.

Statistics of bulletins issued by RSMC New Delhi, Area Cyclone Warning Centre Kolkata, Cyclone Warning Centre Bhubaneswar & Visakhapatnam in association with the SuCS AMPHAN are given in **Table 8**.

**Table 8 (a): Bulletins issued by RSMC New Delhi**

S.N	Bulletin type	No. of Bulletins	Issued to
1	Informatory Message	3	<ol style="list-style-type: none"> <li>1. IMD website, RSMC New Delhi website and Mausam website</li> <li>2. FAX and e-mail to Control Room Ministry of Home Affairs &amp; National Disaster Management Authority, Cabinet Secretariat, Minister of Science &amp; Technology, Headquarter Integrated Defence Staff, Director General Doordarshan, All India Radio, National Disaster Response Force, Press Information Bureau, Chief Secretary to Government of Kerala, Tamil Nadu, Andaman &amp; Nicobar Islands, Andhra Pradesh, Odisha, West Bengal, Assam, Meghalaya, Arunachal Pradesh.</li> <li>3. First informatory message was issued on 13<sup>th</sup> May.</li> </ol>
2	National Bulletin	45	<ol style="list-style-type: none"> <li>1. IMD website, RSMC New Delhi website and Mausam website</li> <li>2. FAX and e-mail to Control Room Ministry of Home Affairs &amp; National Disaster Management Authority, Cabinet Secretariat, Minister of Science &amp; Technology, Headquarter Integrated Defence Staff, Director General Doordarshan, All India Radio, National Disaster Response Force, Press Information Bureau, Chief Secretary to Government of Kerala, Tamil Nadu, Andaman &amp; Nicobar Islands, Andhra Pradesh, Odisha, West Bengal, Assam, Meghalaya, Arunachal Pradesh.</li> <li>3. First Bulletin was issued on 16<sup>th</sup> May morning.</li> </ol>
3	RSMC Bulletin	45	<ol style="list-style-type: none"> <li>1. IMD's website, RSMC website and Mausam website</li> <li>2. WMO/ESCAP member countries including Bangladesh and Myanmar through GTS and E-mail.</li> <li>3. First message was issued on 16<sup>th</sup> May morning.</li> </ol>
4	GMDSS Bulletins	33	<ol style="list-style-type: none"> <li>1. IMD website, RSMC New Delhi website</li> <li>2. Transmitted through WMO Information System (WIS) to Joint WMO/IOC Technical Commission for Ocean and Marine Meteorology (JCOMM)</li> <li>3. First Bulletin was issued on 16<sup>th</sup> May morning.</li> </ol>
5	Tropical Cyclone Advisory Centre Bulletin	22	<ol style="list-style-type: none"> <li>1. Met Watch offices in Asia Pacific regions and middle east through GTS to issue Significant Meteorological information for International Civil Aviation</li> <li>2. WMO's Aviation Disaster Risk Reduction (ADRR), Hong Kong through ftp</li> <li>3. RSMC website</li> <li>4. First message was issued on 16<sup>th</sup> May morning.</li> </ol>
6	Tropical Cyclone Vital Statistics	22	Modelling group of IMD, National Centre for Medium Range Weather Forecasting Centre (NCMRWF), Indian National Centre for Ocean Information Services (INCOIS), Indian Institute of Technology (IIT) Delhi, IIT Bhubaneswar etc.

7	Warnings through SMS	Frequently	<p>SMS to disaster managers at national level and concerned states (every time when there was change in track, intensity and landfall characteristics)</p> <p>(i) 3,14,016 SMS to General Public by IMD Headquarters to users registered at RSMC website <a href="http://www.rsmcnewdelhi.imd.gov.in">www.rsmcnewdelhi.imd.gov.in</a></p> <p>(ii) 2,420 SMS to senior level disaster managers at centre and affected states along the east coast by IMD Headquarters</p> <p>(iii) 75,06,968 SMS to registered users including fishermen by INCOIS</p> <p>(iv) INCOIS also issued 39 INCOIS-IMD joint bulletins (in short template) issued through NAVIC (for communication to fishermen in open sea)</p> <p>(v) 28,78,069 SMS to farmers in the affected regions of Andhra Pradesh, Odisha, West Bengal, Meghalaya and Assam by Kisan Portal</p> <p>(vi) 1,43,41,131 Notifications sent through Mobile App UMANG on 19th and 20th May</p> <p>(vii) Google also issued Cyclone Alert notifications to the people in affected areas of West Bengal and Odisha</p> <p>(viii) Concerned State Governments of Odisha and West Bengal also disseminated warnings using various Mobile communicating systems</p>
8	Warnings through Social Media	Daily	Cyclone Warnings were uploaded on Social networking sites (Face book, Tweeter and WhatsApp) since inception to weakening of system (every time when there was change in track, intensity and landfall characteristics) and hourly on the day of landfall on 20 <sup>th</sup> May.
9	Press conference	Daily	3 – Joint press conferences were held by DG IMD & DG NDRF from National Media Centre on 19 <sup>th</sup> , 20 <sup>th</sup> & 21 <sup>st</sup> May.
10	Press Release	11	Disaster Managers, Media persons by email and uploaded on website
11	Press Briefings	Daily	Regular briefing daily and frequently as and when media persons visited the National Weather Forecasting Centre
12	Hourly Updates	19	Hourly bulletins by email, website, social media

**Table 8(b): Statistics of bulletins issued by ACWC Kolkata and CWC Bhubaneswar & Visakhapatnam**

S.No.	Type of Bulletin	No. of Bulletins issued		
		ACWC Kolkata	CWC Bhubaneswar	CWC Visakhapatnam
1.	Sea Area Bulletin	38	-	-
2.	Coastal Weather Bulletins	38 each for West Bengal and Andaman & Nicobar Islands	29	25
3.	Fishermen Warnings issued	31 each for West Bengal and 32 Andaman & Nicobar Islands	37	35
4.	Port Warnings	31 Hooghly Port & Sagar Islands and 29 Port Blair	36	15
5.	Heavy Rainfall warning	33	09	NIL
6.	Gale Wind Warning	31	36	5
7.	Storm Surge Warning	16	NIL	NIL
8.	Information & Warning issued to State Government and other Agencies	35 Bulletins 500 Phone calls	39 bulletins	10
9.	SMS	SMS NIL Whatsapp-70,000	SMS NIL Whatsapp-41327	SMS 1591 Whatsapp-38 to 55 media members
10.	Press Conference/Briefing/All India Radio	4/50/10	05	07

**7. Major challenges during monitoring and prediction of SuCS AMPHAN:**

There were 4 main challenges while monitoring Amphan.

- i. Amphan moved very slowly during initial 2 days with a speed of 4-5 kmph and very fast during last 2 days prior to landfall with windspeed of about 20-30 kmph. Thus, the movement of the system was not uniform. It was very challenging to determine the speed of movement in different phases accurately and thus determining the landfall time correctly.
- ii. The challenge was more severe while considering the numerical model guidance about the possible track of the cyclone. We usually examine about 12 global and regional models including six models run by Ministry of Earth Sciences and six international models. The model guidance wrt track was

highly inconsistent with variation from day to day and also from morning to evening. There was a large spread in the tracks suggested by different models even two days before the landfall. So developing a consensus based on these models was very challenging.

- iii. Amphan underwent rapid intensification (increase in intensity by 30 knots [55 kmph) in 24 hours] from 17<sup>th</sup> noon (1130 hrs IST) to 19<sup>th</sup> early morning (2100 UTC of 18<sup>th</sup>) which is one of the rare record of intensification in this ocean basin, an increase in wind speed by almost 2.3 times).
- iv. Amphan originated from a low pressure area (LPA) which developed over south Andaman Sea on 1<sup>st</sup> May. It persisted over the same region for about 5 days and became less marked. However, the remnant cyclonic circulation moved gradually west-northwestwards and remained over southeast BoB and adjoining south Andaman Sea for a long time upto 12<sup>th</sup> May. It again reappeared as an LPA on 13<sup>th</sup> May over southeast BoB. Considering the model guidance about genesis (formation of depression- a cyclonic circulation system with wind speed of 32-50 kmph), there was false alarm from 25<sup>th</sup> April onwards about the genesis of the cyclone over the BoB and it's landfall over different coasts like Bangladesh, Myanmar and Andaman & Nicobar Islands. It was a challenge to predict the place and occurrence of LPA and it's possible intensification into a depression, it's intensification and movement towards a particular coast. However, IMD utilized all the objective tools to compare, comprehend and analyse the NWP model guidance and the observations from various satellites, buoys, ships and coastal observations to correctly predict the genesis of LPA on 13<sup>th</sup> May. Also with the formation of LPA on 13<sup>th</sup> May, IMD issued the first Press Release and Special Informatory Message to the concerned states and central government agencies about it's possible intensification into a cyclone and it's movement towards north BoB following a recurving track.

## **8. Initiatives during SuCS AMPHAN:**

- (i) For the first time, during Amphan track was uploaded on GIS based platform
- (ii) Notifications on Amphan were also sent through Govt. of India Mobile App, Umang
- (iii) Notifications were also sent through Google Network.

## **9. Appreciations earned for accurate forecast of SuCS AMPHAN**

- (i) Chief Minister of Odisha appreciated the cyclone warning services of IMD during cyclone Amphan
- (ii) DG NDRF appreciated the accurate predictions of landfall time, point, track and intensity of Amphan that helped them plan evacuation measures effectively

- (iii) Chief Disaster Risk Reduction Centre, UNDP appreciated the accurate warnings including adverse weather warnings during Amphan
- (iv) WMO appreciated the services of RSMC New Delhi during Amphan
- (v) Indian Air Force appreciated the accurate prediction of Amphan by IMD, stating that it gave them ample advance notice to initiate steps to mitigate the damages and casualties.
- (vi) Leading national & international print and electronic media published the story of monitoring of Amphan and life profile of DGM IMD

**10. Acknowledgement:**

India Meteorological Department (IMD) and RSMC New Delhi duly acknowledge the contribution from all the stake holders and disaster management agencies who contributed to the successful monitoring, prediction and early warning service of SuCS AMPHAN. We acknowledge the contribution of all sister organisations of Ministry of Earth Sciences including National Centre for Medium Range Weather Forecasting Centre (NCMRWF), Indian National Centre for Ocean Information Services (INCOIS), National Institute of Ocean Technology (NIOT), Indian Institute of Tropical Meteorology (IITM) Pune, DRDO Integrated Test Range, Chandipur, research institutes including IIT Bhubaneswar, IIT Delhi and Space Application Centre, Indian Space Research Organisation (SAC-ISRO) for their valuable support. The support from various Divisions/Sections of IMD including Area Cyclone Warning Centre (ACWC) Chennai, Kolkata, Cyclone Warning Centre (CWC) Bhubaneswar, Visakhapatnam, Meteorological Centre (MC) Agartala, Doppler Weather Radar Stations at Visakhapatnam, Chandipur, Gopalpur, Paradip, Kolkata & Agartala and coastal observatories of Odisha & north Andhra Pradesh. The contribution from Numerical Weather Prediction Division, Satellite and Radar Divisions, Surface & Upper air instruments Divisions, New Delhi and Information System and Services Division at IMD is also duly acknowledged.

-----

Typical INSAT-3D imageries during life cycle of SUCS APHAN (16<sup>th</sup>-21<sup>st</sup> May) are presented in Fig. 1-5.




Fig. 1: INSAT-3D enhanced colour imageries during life cycle of SuCS AMPHAN (15-21 May, 2020)







Fig. 2: INSAT-3D BD imageries during life cycle of SuCS AMPHAN (15-22 May, 2020)



**Fig. 3: INSAT-3D Visible imageries during life cycle of SuCS AMPHAN (15-22 May, 2020)**




**Fig. 4: INSAT-3D IR imageries during life cycle of SuCS AMPHAN (15-22 May, 2020)**



**Fig. 5: INSAT-3D Cloud Top Brightness Temperature (CTBT) imageries during life cycle of SuCS AMPHAN (15-22 May, 2020)**

Typical SCAT SAT imageries during life cycle of SUCS Amphan (13<sup>th</sup> May-20<sup>th</sup> May) since inception as low pressure area are presented in Fig. 6.



**Fig. 6: SCAT SAT imageries during life cycle of SuCS AMPHAN (13-20 May, 2020)**

Typical ASCAT imageries during life cycle of SUCS Amphan (13<sup>th</sup> -20<sup>th</sup> May), since inception as low pressure area are presented in Fig.7.





Fig. 7: ASCAT imageries during life cycle of SuCS AMPHAN (13-20 May, 2020)

SUCS Amphan was continuously monitored by IMD's Doppler Weather Radars at Visakhapatnam, Gopalpur, Paradeep, Chandipur, Kolkata and Agartala while moving from westcentral BoB to northeast India. Typical radar imageries from Visakhapatnam, Gopalpur, Paradeep, Chandipur and Kolkata radar centres are presented in Fig. 8-12.



**Fig. 8:** Typical Radar Max dBZ imageries from DWR Visakhapatnam during 18-19 May, 2020






Fig. 10 a: Typical Radar Max Z imageries of DWR Paradeep during during 19-20 May, 2020






Fig. 10 b: Typical Radar Max Z imageries of DWR Paradeep during during 20-21 May, 2020




**Fig. 11:** Typical Radar MaxZ imageries of 'DRDO Integrated Test Range', Chandipur during 19-20 May, 2020




**Fig. 12:** Typical Radar Max dBZ imageries from DWR Kolkata during 19-20 May, 2020

The total precipitable water (TPW) imageries (Source: TC Forecaster Website: [https://rammb-data.cira.colostate.edu/tc\\_realtime/index.asp](https://rammb-data.cira.colostate.edu/tc_realtime/index.asp)) during life cycle of SUCS Amphan are presented in Fig. 13. These imageries indicate excessive increase in warm moist around the system centre on 1330 UTC of 16<sup>th</sup> May. The warm moist air supply continued till 1800 UTC of 18<sup>th</sup> May. Thereafter, it gradually decreased. The system also exhibited decrease in intensity on 19<sup>th</sup> & 20<sup>th</sup> prior to landfall.




**Fig. 13: Typical total precipitable water imageries during life cycle of SuCS Amphan (16<sup>th</sup>-21<sup>st</sup> May, 2020).**

IMD GFS analysis of mean sea level pressure, winds at 10m, 850 hPa, 500 hPa and 200 hPa levels based on 0000 UTC during 13<sup>th</sup> -21<sup>st</sup> May, 2020 are presented in Fig.14. On 13<sup>th</sup> May, IMD GFS indicated a low pressure area over southeast BOB with vertical extension of the cyclonic circulation upto 500 hPa level.




**Fig. 14 (a):** IMD GFS (T574) mean sea level pressure (MSLP), winds at 10m, 850, 500 and 200 hPa levels based on 0000 UTC of 13<sup>th</sup> May,2020

On 14<sup>th</sup> May, IMD GFS indicated a low pressure area over southwest & adjoining southeast BOB with vertical extension of the cyclonic circulation upto 500 hPa level.




**Fig. 14 (b): IMD GFS (T574) mean sea level pressure (MSLP), winds at 10m, 850, 500 and 200 hPa levels based on 0000 UTC of 14<sup>th</sup> May,2020**

On 15<sup>th</sup> May, IMD GFS indicated a well marked low pressure area over southwest BOB with vertical extension of the cyclonic circulation upto 200 hPa level.




**Fig. 14 (c): IMD GFS (T574) mean sea level pressure (MSLP), winds at 10m, 850, 500 and 200 hPa levels based on 0000 UTC of 15<sup>th</sup> May, 2020**

On 16<sup>th</sup> May, IMD GFS indicated a deep depression over southwest BOB with vertical extension of the cyclonic circulation upto 500 hPa level. Actually, it was a depression on 16<sup>th</sup> May over southeast BoB. IMD GFS slightly over estimated the intensity of the system.




**Fig. 14 (d): IMD GFS (T574) mean sea level pressure (MSLP), winds at 10m, 850, 500 and 200 hPa levels based on 0000 UTC of 16<sup>th</sup> May,2020**

On 17<sup>th</sup> May, IMD GFS indicated rapid intensification of the system It lay as a severe cyclonic storm over southwest BOB with vertical extension of the cyclonic circulation upto 200 hPa level. GFS also indicated near northwards movement of the system. Actually, it was a cyclonic storm at 0000 UTC of 17<sup>th</sup> May over southeast BoB. IMD GFS over estimated the intensity of the system.




**Fig. 14 (e): IMD GFS (T574) mean sea level pressure (MSLP), winds at 10m, 850, 500 and 200 hPa levels based on 0000 UTC of 17 May,2020**

On 18<sup>th</sup> May, IMD GFS indicated further intensification of the system. It lay as a very severe cyclonic storm over westcentral BOB with vertical extension of the cyclonic circulation upto 200 hPa level. GFS also indicated near northwards movement of the system. Actually, it was an extremely severe cyclonic storm at 0000 UTC of 18<sup>th</sup> May over westcentral BoB. IMD GFS correctly picked up intensity and movement of the system.




**Fig. 14 (f): IMD GFS (T574) mean sea level pressure (MSLP), winds at 10m, 850, 500 and 200 hPa levels based on 0000 UTC of 18<sup>th</sup> May,2020**

On 19<sup>th</sup> May, IMD GFS indicated a very intense system over westcentral BOB with vertical extension of the cyclonic circulation upto 200 hPa level. GFS also indicated northeastwards movement of the system. Actually, it was a super cyclonic at 0000 UTC of 19<sup>th</sup> May over westcentral BoB. IMD GFS correctly picked up intensity and movement of the system.




**Fig.14 (g): IMD GFS (T574) mean sea level pressure (MSLP), winds at 10m, 850, 500 and 200 hPa levels based on 0000 UTC of 19<sup>th</sup> May,2020**

On 20<sup>th</sup> May, IMD GFS indicated slight weakening of the system. It lay as a very severe cyclonic storm over northwest BoB off Odisha coast. The system extended vertically upto 200 hPa level. GFS also indicated northeastwards movement of the system. Actually, it was an extremely severe cyclonic storm cyclonic storm at 0000 UTC of 20<sup>th</sup> May over northwest BoB. GFS also picked up an anticyclone over Myanmar to the west of system and strong westerlies in the upper level indicating steering of the system in northeastwards direction. IMD GFS correctly picked up intensity and movement of the system.



**Fig. 14 (h): IMD GFS (T574) mean sea level pressure (MSLP), winds at 10m, 850, 500 and 200 hPa levels based on 0000 UTC of 20<sup>th</sup> May, 2020**

On 21<sup>st</sup> May, IMD GFS indicated a cyclonic storm over Bangladesh and adjoining Gangetic West Bengal.



**Fig. 14 (i): IMD GFS (T574) mean sea level pressure (MSLP), winds at 10m, 850, 500 and 200 hPa levels based on 0000 UTC of 21<sup>st</sup> May,2020**

IMD GFS thus correctly picked up intensification, movement and weakening of the system.